

**Diptera Greenland collection of the Natural History Museum of Denmark (Zoological Museum),
University of Copenhagen**

as per September 4th 2002.

The collection of Diptera from Greenland includes alcohol and dry material. The following list is complete only for the dry material and the families are in alphabetic order.

The collection contains about 22,000 specimens in 250 species from 30 families.

At the end of the document, you will find a list of our Greenland material that is not sorted to species

| Family | Genus | Species | Author | Number of specimens |
|---------------|-----------------------|----------------------|---------------------------|----------------------------|
| Agromyzidae | <i>Cerodontha</i> | <i>lindrothi</i> | Griffiths, 1964 | 3 |
| Agromyzidae | <i>Cerodontha</i> | <i>thulensis</i> | Griffiths, 1966 | 3 |
| Agromyzidae | <i>Napomyza</i> | <i>parvicella</i> | Coquillett, 1902) | 3 |
| Agromyzidae | <i>Phytoliriomyza</i> | <i>arctica</i> | (Lundbeck, 1901) | 104 |
| Agromyzidae | <i>Phytomyza</i> | <i>aquilonia</i> | Frey, 1946 | 12 |
| Agromyzidae | <i>Phytomyza</i> | <i>fuscula</i> | Zetterstedt, 1838 | 87 |
| Agromyzidae | <i>Phytomyza</i> | <i>opacella</i> | Hendel, 1935 | 12 |
| Agromyzidae | <i>Phytomyza</i> | <i>ranunculi</i> | (Schrank, 1803) | 16 |
| Agromyzidae | <i>Phytomyza</i> | <i>varipes</i> | Macquart, 1835 | 2 |
| Anthomyiidae | <i>Botanophila</i> | <i>betarum</i> | (Lintner, 1883) | 104 |
| Anthomyiidae | <i>Botanophila</i> | <i>moriens</i> | (Zetterstedt, 1845) | 3 |
| Anthomyiidae | <i>Botanophila</i> | <i>profuga</i> | (Stein, 1916) | 66 |
| Anthomyiidae | <i>Botanophila</i> | <i>rubrigena</i> | (Schnabl, 1915) | 54 |
| Anthomyiidae | <i>Delia</i> | <i>echinata</i> | (Séguy, 1923) | 129 |
| Anthomyiidae | <i>Delia</i> | <i>fabricii</i> | (Homgren, 1873) | 143 |
| Anthomyiidae | <i>Delia</i> | <i>groenlandica</i> | Griffiths, 1993 | 13 |
| Anthomyiidae | <i>Delia</i> | <i>platura</i> | (Meigen, 1826) | 38 |
| Anthomyiidae | <i>Delia</i> | <i>radicum</i> | (Linnaeus, 1758) | 15 |
| Anthomyiidae | <i>Egle</i> | <i>bicaudata?</i> | (Malloch, 1920) | 80 |
| Anthomyiidae | <i>Eutrichota</i> | <i>tunicata</i> | (Zetterstedt, 1846) | 6 |
| Anthomyiidae | <i>Fucellia</i> | <i>ariciiformis</i> | (Holmgren, 1872) | 84 |
| Anthomyiidae | <i>Fucellia</i> | <i>fucorum</i> | (Fallén, 1819) | 102 |
| Anthomyiidae | <i>Fucellia</i> | <i>maritima</i> | (Haliday, 1838) | 11 |
| Anthomyiidae | <i>Fucellia</i> | <i>pictipennis</i> | Becker, 1907 | 26 |
| Anthomyiidae | <i>Fucellia</i> | <i>tergina</i> | (Zetterstedt, 1845) | 6 |
| Anthomyiidae | <i>Fucellia</i> | <i>vibei</i> | Collin, 1951 | 2 |
| Anthomyiidae | <i>Lasiomma</i> | <i>octoguttatum</i> | (Zetterstedt, 1845) | 32 |
| Anthomyiidae | <i>Lasiomma</i> | <i>picipes</i> | (Meigen, 1826) | 218 |
| Anthomyiidae | <i>Paradelia</i> | <i>intersecta</i> | (Meigen, 1826) | 26 |
| Anthomyiidae | <i>Paregle</i> | <i>audacula</i> | (Harris, 1780) | 32 |
| Anthomyiidae | <i>Pegohylemyia</i> | <i>biseriata</i> | Huckett | 1 |
| Anthomyiidae | <i>Pegomya</i> | <i>circumpolaris</i> | Ackland & Griffiths, 1983 | 20 |
| Anthomyiidae | <i>Pegomya</i> | <i>flavifrons</i> | (Walker, 1849) | 6 |
| Anthomyiidae | <i>Pegomya</i> | <i>ruficeps</i> | Fallén | 1 |
| Anthomyiidae | <i>Pegomya</i> | <i>tenera</i> | (Zetterstedt, 1838) | 16 |
| Anthomyiidae | <i>Pegomya</i> | <i>zonata</i> | (Zetterstedt, 1838) | 32 |

| | | | | |
|---------------|------------------------|------------------------|---------------------------|-----|
| Anthomyiidae | <i>Pegomyia</i> | <i>circumpolaris</i> | Ackland & Griffiths, 1983 | 16 |
| Anthomyiidae | <i>Pegomyia</i> | <i>flavifrons</i> | (Walker, 1849) | 13 |
| Anthomyiidae | <i>Pegomyia</i> | <i>icterica</i> | (Holmgren, 1873) | 22 |
| Anthomyiidae | <i>Pegomyia</i> | <i>notabilis</i> | (Zetterstedt, 1846) | 77 |
| Anthomyiidae | <i>Pegomyia</i> | <i>zonata</i> | (Zetterstedt, 1838) | 20 |
| Anthomyiidae | <i>Pegoplata</i> | <i>tundrica</i> | (Schnabl, 1915) | 104 |
| Anthomyiidae | <i>Zaphne</i> | <i>bruneifrons</i> | (Zetterstedt, 1838) | 13 |
| Anthomyiidae | <i>Zaphne</i> | <i>divisa</i> | (Meigen, 1826) | 7 |
| Anthomyiidae | <i>Zaphne</i> | <i>frontata</i> | (Zetterstedt, 1838) | 106 |
| Anthomyiidae | <i>Zaphne</i> | <i>nearctica</i> | (Huckett, 1965) | 8 |
| Anthomyiidae | <i>Zaphne</i> | <i>occidentalis</i> | (Malloch, 1920) | 5 |
| Anthomyiidae | <i>Zaphne</i> | <i>verticina</i> | (Zetterstedt, 1838) | 29 |
| Calliphoridae | <i>Boreellus</i> | <i>atriceps</i> | (Zetterstedt, 1845) | 45 |
| Calliphoridae | <i>Calliphora</i> | <i>uralensis</i> | Villeneuve, 1922 | 85 |
| Calliphoridae | <i>Calliphora</i> | <i>vicina</i> | Robineau-Desvoidy, 1830 | 34 |
| Calliphoridae | <i>Cynomyia</i> | <i>mortuorum</i> | (Linnaeus, 1761) | 139 |
| Calliphoridae | <i>Phormia</i> | <i>terraenovae</i> | Robineau-Desvoidy, 1830 | 329 |
| Calliphoridae | <i>Protocalliphora</i> | <i>azurea</i> | (Fallén, 1817) | 3 |
| Cecidomyiidae | <i>Xylopriona</i> | <i>atra</i> | (Meigen, 1804) | 3 |
| Chironomidae | <i>Ablabesmyia</i> | <i>pulchripennis</i> | (Lundbeck, 1898) | 38 |
| Chironomidae | <i>Anatopynia</i> | <i>pictipennis</i> | (Zetterstedt, 1838) | 1 |
| Chironomidae | <i>Chironomus</i> | <i>hyperboreus</i> | Staeger, 1845 | 3 |
| Chironomidae | <i>Chironomus</i> | n. sp. | | 8 |
| Chironomidae | <i>Chironomus</i> | <i>riparius</i> | Meigen, 1804 | 18 |
| Chironomidae | <i>Chironomus</i> | <i>staegeri</i> | Lundbeck, 1898 | 17 |
| Chironomidae | <i>Corynoneura</i> | <i>scutellata</i> | Winnertz, 1846) | 33 |
| Chironomidae | <i>Cricotopus</i> | <i>glacialis</i> | Edwards, 1922 | 5 |
| Chironomidae | <i>Cricotopus</i> | <i>tibialis</i> | (Meigen, 1804) | 1 |
| Chironomidae | <i>Diamesa</i> | <i>aberrata?</i> | Lundbeck, 1889 | 8 |
| Chironomidae | <i>Diamesa</i> | <i>biappendiculata</i> | Goetghebuer in Remy, 1928 | 1 |
| Chironomidae | <i>Diamesa</i> | <i>gregsoni</i> | Edwards, 1933 | 12 |
| Chironomidae | <i>Dicrotendipes</i> | <i>nervosus</i> | (Staeger, 1839) | 1 |
| Chironomidae | <i>Eukiefferiella</i> | <i>claripennis</i> | (Lundbeck, 1898) | 5 |
| Chironomidae | <i>Halocladus</i> | <i>vitripennis</i> | (Meigen, 1818) | 38 |
| Chironomidae | <i>Limnophyes</i> | <i>atomarius</i> | (Zetterstedt, 1850) | 2 |
| Chironomidae | <i>Limnophyes</i> | <i>borealis</i> | Goetghebuer, 1933 | 1 |
| Chironomidae | <i>Limnophyes</i> | cf. <i>minima</i> | (Meigen, 1818) | 1 |
| Chironomidae | <i>Limnophyes</i> | <i>eltoni</i> | (Edwards, 1922) | 1 |
| Chironomidae | <i>Limnophyes</i> | <i>pumillio</i> | (Holmgren, 1869) | 48 |
| Chironomidae | <i>Metriocnemus</i> | <i>atratus</i> | (Zetterstedt, 1850) | 6 |
| Chironomidae | <i>Metriocnemus</i> | <i>debilipennis</i> | (Lundbeck, 1898) | 1 |
| Chironomidae | <i>Metriocnemus</i> | <i>fuscipes</i> | (Meigen, 1818) | 20 |
| Chironomidae | <i>Metriocnemus</i> | <i>lundbeckii</i> | Johannsen, 1905 | |
| Chironomidae | <i>Metriocnemus</i> | <i>similis</i> | Kieffer, 1922 | 1 |
| Chironomidae | <i>Metriocnemus</i> | <i>ursinus</i> | (Holmgren, 1869) | 11 |

| | | | | |
|----------------|-----------------------|----------------------------|----------------------------|-----|
| Chironomidae | <i>Micropsectra</i> | <i>groenlandica</i> | Andersen, 1937 | 2 |
| Chironomidae | <i>Micropsectra</i> | <i>junci</i> | (Meigen, 1818) | 3 |
| Chironomidae | <i>Orthocladius</i> | <i>decoratus</i> | (Holmgren, 1869) | 1 |
| Chironomidae | <i>Orthocladius</i> | <i>difficilis</i> | (Lundbeck, 1898) | 13 |
| Chironomidae | <i>Orthocladius</i> | <i>frigidus</i> | (Zetterstedt, 1838) | 20 |
| Chironomidae | <i>Orthocladius</i> | <i>graminicola</i> | (Lundbeck, 1898) | 21 |
| Chironomidae | <i>Orthocladius</i> | <i>minutus</i> | (Zetterstedt, 1850) | 17 |
| Chironomidae | <i>Paratanytarsus</i> | <i>setosimanus</i> | (Goetghebuer, 1928) | 46 |
| Chironomidae | <i>Paratanytarsus</i> | <i>tenuis</i> | (Meigen, 1839) | 2 |
| Chironomidae | <i>Parochlus</i> | <i>kiefferi</i> | (Garret 1925) | 1 |
| Chironomidae | <i>Podonomus</i> | <i>posticalis</i> | Lundbeck | 4 |
| Chironomidae | <i>Procladius</i> | <i>choreus</i> | (Meigen, 1804) | 32 |
| Chironomidae | <i>Procladius</i> | <i>crassinervis</i> | (Zetterstedt, 1838) | 3 |
| Chironomidae | <i>Psectrocladius</i> | <i>armatus</i> | Andersen | 2 |
| Chironomidae | <i>Psectrocladius</i> | <i>limbatellus</i> | (Holmgren, 1869) | 40 |
| Chironomidae | <i>Psilodiamesa</i> | <i>abelsini</i> | Vibe | 12 |
| Chironomidae | <i>Psilodiamesa</i> | <i>chorea</i> | Lundbeck, 1898 | 26 |
| Chironomidae | <i>Sergentia</i> | <i>coracina</i> | (Zetterstedt, 1850) | 10 |
| Chironomidae | <i>Smittia</i> | <i>extrema</i> | (Holmgren, 1869) | 22 |
| Chironomidae | <i>Smittia</i> | <i>longitibia</i> | (Goetghebuer, 1933) | 7 |
| Chironomidae | <i>Smittia</i> | <i>parva</i> | (Lundbeck, 1898) | 8 |
| Chironomidae | <i>Smittia</i> | <i>polymorpha</i> | Andersen, 1937 | 12 |
| Chironomidae | <i>Smittia</i> | <i>subnudipennis</i> | (Goetghebuer, 1933) | 4 |
| Chironomidae | <i>Smittia</i> | <i>velutina</i> | (Lundbeck, 1898) | 6 |
| Chironomidae | <i>Tanytarsus</i> | <i>anderseni</i> | Reiss & Pittkau | 5 |
| Chironomidae | <i>Tanytarsus</i> | <i>niger</i> | Andersen, 1937 | 3 |
| Chironomidae | <i>Tendipes</i> | <i>dorsalis riparius</i> | various Authors | 1 |
| Chironomidae | <i>Tendipes</i> | <i>hyperborea</i> | Staeger, 1845 | 13 |
| Chironomidae | <i>Tendipes</i> | <i>staegeri</i> | Lundbeck, 1898 | 9 |
| Chironomidae | <i>Trichocladius</i> | <i>chapmani</i> | (Edwards, 1935) | 3 |
| Chironomidae | <i>Trichocladius</i> | <i>pubitarsis</i> | (Zetterstedt, 1838) | 1 |
| Culicidae | <i>Aedes</i> | <i>impiger</i> | (Walker, 1848) | 13 |
| Culicidae | <i>Aedes</i> | <i>nigripes</i> | (Zetterstedt, 1838) | 15 |
| Dolichopodidae | <i>Dolichopus</i> | <i>dasyops</i> | Malloch, 1919 | 19 |
| Dolichopodidae | <i>Dolichopus</i> | <i>groenlandicus</i> | Zetterstedt, 1843 | 540 |
| Dolichopodidae | <i>Dolichopus</i> | <i>humilis</i> | Van Duzee, 1921 | 51 |
| Dolichopodidae | <i>Dolichopus</i> | <i>plumipes</i> | (Scopoli, 1763) | 86 |
| Empididae | <i>Clinocera</i> | <i>stagnalis</i> | (Haliday, 1833) | 40 |
| Empididae | <i>Rhamphomyia</i> | <i>filicauda</i> | Henriksen & Lundbeck, 1917 | 34 |
| Empididae | <i>Rhamphomyia</i> | <i>hirticula</i> | Collin, 1937 | 2 |
| Empididae | <i>Rhamphomyia</i> | <i>hirtula</i> | Zetterstedt, 1842 | 158 |
| Empididae | <i>Rhamphomyia</i> | <i>hoeli</i> | Frey, 1950 | 33 |
| Empididae | <i>Rhamphomyia</i> | <i>nigrita</i> | (Zetterstedt, 1838) | 240 |
| Ephydridae | <i>Lamproscatella</i> | <i>brunnipes norvegica</i> | Andersson, 1975 | 50 |
| Ephydridae | <i>Lamproscatella</i> | <i>quadrisetosa</i> | (Becker, 1896) | 23 |

| | | | | |
|--------------|-------------------------|-----------------------|---------------------------|-----|
| Ephydridae | <i>Limnellia</i> | <i>stenhammeri</i> | (Zetterstedt, 1846) | 25 |
| Ephydridae | <i>Napaea</i> | <i>pusilla</i> | (Miegen, 1830) | 1 |
| Ephydridae | <i>Philygria</i> | <i>vittipennis</i> | (Zetterstedt, 1838) | 5 |
| Ephydridae | <i>Scatella</i> | <i>stagnalis</i> | (Fallén, 1823) | 137 |
| Ephydridae | <i>Scatella</i> | <i>tenuicosta</i> | Collin, 1930 | 112 |
| Ephydridae | <i>Scatophila</i> | <i>cribata</i> | (Stenhammar, 1844) | 26 |
| Fannidae | <i>Fannia</i> | <i>bifimbriata</i> | Collin in Carpenter, 1938 | 4 |
| Fannidae | <i>Fannia</i> | <i>canicularis</i> | (Linnaeus, 1761) | 33 |
| Fannidae | <i>Fannia</i> | <i>scalaris</i> | (Fabricius, 1794) | 10 |
| Heleomyzidae | <i>Heleomyza</i> | <i>borealis</i> | (Boheman, 1865) | 151 |
| Heleomyzidae | <i>Neoleria</i> | <i>prominens</i> | (Becker, 1897) | 72 |
| Heleomyzidae | <i>Scoliocentra</i> | <i>fraterna?</i> | Loew, 1863 | 53 |
| Limoniidae | <i>Ormosia</i> | <i>affinis</i> | (Lundbeck, 1898) | 30 |
| Limoniidae | <i>Ormosia</i> | <i>fascipennis</i> | (Zetterstedt, 1838) | 72 |
| Milichiidae | <i>Desmometopa</i> | <i>sordida?</i> | (Fallén, 1820) | 1 |
| Muscidae | ? | <i>brunneifrons?</i> | | 83 |
| Muscidae | ? | <i>tundrica</i> | | 45 |
| Muscidae | <i>Drymeia</i> | <i>segnis</i> | (Holmgren, 1883) | 112 |
| Muscidae | <i>Eupogonomyia</i> | <i>groenlandica</i> | (Lundbeck, 1901) | 74 |
| Muscidae | <i>Graphomya</i> | <i>ungava</i> | Arntfield, 1975 | 14 |
| Muscidae | <i>Hydrophoria</i> | <i>frontata</i> | (Zetterstedt, 1838) | 330 |
| Muscidae | <i>Hydrotaea</i> | <i>anxia</i> | (Zetterstedt, 1838) | 111 |
| Muscidae | <i>Limnophora</i> | <i>sinuata</i> | Collin, 1930 | 170 |
| Muscidae | <i>Lispe</i> | <i>uliginosa</i> | Fallén, 1825 | 35 |
| Muscidae | <i>Lophosceles</i> | <i>frenatus</i> | (Holmgren, 1872) | 85 |
| Muscidae | <i>Musca</i> | <i>domestica</i> | Linnaeus, 1758 | 2 |
| Muscidae | <i>Phaonia</i> | <i>bidentata</i> | (Ringdahl, 1933) | 63 |
| Muscidae | <i>Phaonia</i> | <i>minima</i> | Malloch, 1919 | 15 |
| Muscidae | <i>Phaonia</i> | <i>pallidisquama</i> | (Zetterstedt, 1849) | 86 |
| Muscidae | <i>Pseudolimnophora</i> | <i>rotundata</i> | (Collin, 1930) | 6 |
| Muscidae | <i>Spilogona</i> | <i>almqvisti</i> | (Holmgren, 1880) | 316 |
| Muscidae | <i>Spilogona</i> | <i>alpica</i> | (Zetterstedt, 1845) | 27 |
| Muscidae | <i>Spilogona</i> | <i>arctica</i> | Huckett, 1965 | 4 |
| Muscidae | <i>Spilogona</i> | <i>baltica</i> | (Ringdahl, 1918) | 12 |
| Muscidae | <i>Spilogona</i> | <i>contractifrons</i> | (Zetterstedt, 1838) | 878 |
| Muscidae | <i>Spilogona</i> | <i>deflorata</i> | (Holmgren, 1872) | 117 |
| Muscidae | <i>Spilogona</i> | <i>denudata</i> | (Holmgren, 1869) | 77 |
| Muscidae | <i>Spilogona</i> | <i>digressa</i> | Huckett, 1965 | 7 |
| Muscidae | <i>Spilogona</i> | <i>dorsata</i> | (Zetterstedt, 1845) | 247 |
| Muscidae | <i>Spilogona</i> | <i>extensa</i> | (Malloch, 1920) | 104 |
| Muscidae | <i>Spilogona</i> | <i>latilamina</i> | (Collin, 1930) | 226 |
| Muscidae | <i>Spilogona</i> | <i>malaisei</i> | (Ringdahl, 1920) | 70 |
| Muscidae | <i>Spilogona</i> | <i>megastoma</i> | (Boheman, 1866) | 97 |
| Muscidae | <i>Spilogona</i> | <i>melanosoma</i> | (Huckett, 1932) | 47 |
| Muscidae | <i>Spilogona</i> | <i>micans</i> | (Ringdahl, 1918) | 23 |

| | | | | |
|----------------|------------------------|------------------------|----------------------------------|-----|
| Muscidae | <i>Spilogona</i> | <i>monacantha</i> | (Collin, 1930) | 29 |
| Muscidae | <i>Spilogona</i> | <i>obsoleta</i> | (Malloch, 1920) | 85 |
| Muscidae | <i>Spilogona</i> | <i>opaca</i> | (Schnabl in Becker et al., 1915) | 29 |
| Muscidae | <i>Spilogona</i> | <i>puberula</i> | (Ringdahl, 1918) | 7 |
| Muscidae | <i>Spilogona</i> | <i>sanctipauli</i> | (Malloch, 1921) | 626 |
| Muscidae | <i>Spilogona</i> | <i>tornensis</i> | (Ringdahl, 1926) | 29 |
| Muscidae | <i>Spilogona</i> | <i>trigonifera</i> | (Zetterstedt, 1838) | 68 |
| Muscidae | <i>Spilogona</i> | <i>tundrae</i> | (Schnabl in Becker et al., 1915) | 147 |
| Mycetophilidae | <i>Boletina</i> | <i>arctica</i> | Holmgren, 1872 | 76 |
| Mycetophilidae | <i>Boletina</i> | <i>sciarina</i> | Staeger, 1840 | 12 |
| Mycetophilidae | <i>Exechia</i> | <i>bicineta</i> | (Staeger, 1840) | 1 |
| Mycetophilidae | <i>Exechia</i> | <i>frigida</i> | (Boheman, 1865) | 1 |
| Mycetophilidae | <i>Gnoriste</i> | <i>apicalis?</i> | Meigen, 1818 | 2 |
| Mycetophilidae | <i>Gnoriste</i> | <i>longirostris</i> | Siebke, 1864 | 2 |
| Mycetophilidae | <i>Gnoriste</i> | <i>longirostris</i> | Siebke, 1864 | 3 |
| Mycetophilidae | <i>Micropsectra</i> | <i>groenlandica</i> | Andersen, 1937 | 32 |
| Mycetophilidae | <i>Mycetophila</i> | <i>fungorum</i> | (De Geer, 1776) | 23 |
| Mycetophilidae | <i>Phronia</i> | <i>exigua</i> | (Zetterstedt, 1852) | 3 |
| Mycetophilidae | <i>Sciophila</i> | <i>hirta</i> | Meigen, 1818 | 3 |
| Mycetophilidae | <i>Trichante</i> | <i>obesa</i> | Winnertz, 1963 | 1 |
| Oestridae | <i>Cephenomyia</i> | <i>trompe</i> | (Modeer, 1786) | 6 |
| Oestridae | <i>Oedemagena</i> | <i>tarandi</i> | (Linnaeus, 1758) | 5 |
| Phoridae | <i>Megaselia</i> | <i>cirriventris</i> | Schmitz, 1929 | 13 |
| Phoridae | <i>Megaselia</i> | <i>clara</i> | (Schmitz, 1921) | 115 |
| Phoridae | <i>Megaselia</i> | <i>groenlandica</i> | (Lundbeck, 1901) | 70 |
| Phoridae | <i>Megaselia</i> | <i>rufipes</i> | (Meigen, 1804) | 14 |
| Piophilidae | <i>Arctopiophila</i> | <i>nigerrima</i> | (Lundbeck, 1901) | 4 |
| Piophilidae | <i>Lasiopiophila</i> | <i>pilosa</i> | (Staeger, 1845) | 47 |
| Piophilidae | <i>Parapiophila</i> | <i>vulgaris</i> | (Fallén, 1820) | 47 |
| Piophilidae | <i>Piophila</i> | <i>casei</i> | (Linnaeus, 1758) | 16 |
| Scathophagidae | <i>Microprosopa</i> | <i>haemorrhoidalis</i> | (Meigen, 1826) | 84 |
| Scathophagidae | <i>Scathophaga</i> | <i>apicalis</i> | (Curtis in Ross, 1835) | 44 |
| Scathophagidae | <i>Scathophaga</i> | <i>litorea</i> | (Fallén, 1819) | 215 |
| Scathophagidae | <i>Scathophaga</i> | <i>nigripalpis</i> | (Becker, 1907) | 4 |
| Scathophagidae | <i>Scatophaga</i> | <i>furcata</i> | (Say, 1823) | 591 |
| Scatopsidae | <i>Scatopse</i> | <i>notata</i> | (Linnaeus, 1758) | 20 |
| Sciaridae | <i>Bradysia</i> | <i>biformis</i> | (Lundbeck, 1898) | 10 |
| Sciaridae | <i>Bradysia</i> | <i>forcipulata</i> | (Lundbeck, 1898) | 12 |
| Sciaridae | <i>Bradysia</i> | <i>iridipennis</i> | (Zetterstedt, 1838) | 2 |
| Sciaridae | <i>Lycoriella</i> | <i>cochleata</i> | (Rübsaamen, 1898) | |
| Sciaridae | <i>Lycoriella</i> | <i>permutata</i> | (Lundbeck, 1900) | |
| Sciaridae | <i>Schwenckfeldina</i> | <i>tridentata</i> | (Rübsaamen, 1898) | 6 |
| Sciaridae | <i>Sciara</i> | <i>?fucata</i> | Meigen | 2 |
| Sciaridae | <i>Sciara</i> | <i>?pallidiventris</i> | Holmg. | 15 |
| Sciaridae | <i>Sciara</i> | <i>aprilina</i> | Meigen, 1818 | 21 |

| | | | | |
|-----------------|-----------------------|----------------------|---------------------------------|------|
| Sciaridae | <i>Sciara</i> | <i>attenuata</i> | Rübsaamen, 1898 | |
| Sciaridae | <i>Sciara</i> | <i>fumatella</i> | Lundbeck, 1898 | |
| Sciaridae | <i>Sciara</i> | <i>humicola</i> | Lundbeck, 1898 | |
| Sciaridae | <i>Sciara</i> | <i>nigripes?</i> | Meigen, 1830 | 3 |
| Sciaridae | <i>Sciara</i> | <i>pulicaria?</i> | Meigen, 1818 | 16 |
| Sepsidae | <i>Orygma</i> | <i>luctuosum</i> | Meigen, 1830 | 21 |
| Sepsidae | <i>Themira</i> | <i>arctica</i> | (Becker in Becker et al., 1915) | 4 |
| Simulidae | <i>Simulium</i> | <i>reptans</i> | (Linnaeus, 1758) | 48 |
| Simulidae | <i>Simulium</i> | <i>vittatum</i> | Zetterstedt, 1838 | 105 |
| Sphaeroceridae | <i>Pteremis</i> | <i>fenestralis?</i> | (Fallén, 1820) | 8 |
| Syrphidae | <i>Carposcalis</i> | <i>lundbecki</i> | (Collin 1931) | 150 |
| Syrphidae | <i>Eupeodes</i> | <i>curtus</i> | (Hine 1922) | 80 |
| Syrphidae | <i>Eupeodes</i> | <i>luniger</i> | (Meigen 1822) | 31 |
| Syrphidae | <i>Eupeodes</i> | <i>nigroventris</i> | (Fluke 1933) | 29 |
| Syrphidae | <i>Helophilus</i> | <i>borealis</i> | Staeger 1845 | 90 |
| Syrphidae | <i>Helophilus</i> | <i>groenlandicus</i> | (Fabricius 1780) | 106 |
| Syrphidae | <i>Neocnemodon</i> | <i>carinatus</i> | (Curran, 1921) | 36 |
| Syrphidae | <i>Parasyrphus</i> | <i>tarsatus</i> | (Zetterstedt 1838) | 240 |
| Syrphidae | <i>Platycheirus</i> | <i>hyperboreus</i> | (Staeger 1845) | 190 |
| Syrphidae | <i>Platycheirus</i> | <i>varipes</i> | Curran 1923 | 52 |
| Syrphidae | <i>Sphaerophoria</i> | <i>scripta</i> | (Linnaeus, 1758) | 225 |
| Syrphidae | <i>Syrphus</i> | <i>torvus</i> | Osten-Sacken, 1875 | 51 |
| Tachinidae | <i>Peleteria</i> | <i>aenea</i> | (Staeger in Zetterstedt, 1849) | 45 |
| Tachinidae | <i>Petinarctia</i> | <i>stylata</i> | (Brauer & Bergenstamm, 1891) | 28 |
| Tachinidae | <i>Trafoia</i> | <i>arctica</i> | (Sack, 1923) | 8 |
| Tethinidae | <i>Pelomyiella</i> | <i>mallochi</i> | (Sturtevant, 1923) | 15 |
| Tipulidae | <i>Erioptera</i> | <i>scotica</i> | Edwards | 23 |
| Tipulidae | <i>Nephrotoma</i> | <i>flavescens</i> | (Linnaeus, 1758) | 1 |
| Tipulidae | <i>Nephrotoma</i> | <i>lundbecki</i> | (Nielsen, 1907) | 28 |
| Tipulidae | <i>Rhabdomastix</i> | <i>caudata</i> | (Lundbeck, 1898) | 2 |
| Tipulidae | <i>Tipula</i> | <i>arctica</i> | Curtis, 1835 | 265 |
| Trichoceridae | <i>Metatrachocera</i> | <i>lutea</i> | Becher, 1886 | 24 |
| Trichoceridae | <i>Trichocera</i> | <i>borealis</i> | Lackschewitz, 1934 | 276 |
| Trichoceridae | <i>Trichocera</i> | <i>maculipennis</i> | Meigen, 1818 | 18 |
| Trichoceridae | <i>Trichocera</i> | <i>regelationis</i> | (Linnaeus, 1758) | 23 |
| Trichoceridae | <i>Trichocera</i> | <i>saltator</i> | (Harris, 1776) | 2 |
| Phoridae | | spp. | | 38 |
| Agromyzidae | | spp. | | 114 |
| Anthomyiidae | <i>Chortophila</i> | spp. | | 6 |
| Anthomyiidae | <i>Fucellia</i> | spp. | | 27 |
| Anthomyiidae | <i>Fucellia</i> | spp. | | 3 |
| Anthomyiidae | | spp. | | 1500 |
| Calliphoridae | | spp. | | 205 |
| Cecidomyiidae | | spp. | | 2 |
| Ceratopogonidae | | spp. | | 10 |

| | | | |
|-----------------------------|----------------------|-------|------|
| Chironomidae | <i>Camptocladius</i> | spp. | 15 |
| Chironomidae | <i>Metriocnemus</i> | spp. | 1 |
| Chironomidae | <i>Metriocnemus</i> | spp. | 1 |
| Chironomidae | <i>Orthocladius</i> | spp. | 57 |
| Chironomidae | <i>Smittia</i> | spp. | 290 |
| Chironomidae | | spp. | 44 |
| Chironomidae | | spp. | 830 |
| Chironomidae | | spp. | 44 |
| Culicidae | | spp. | 450 |
| Dolichopodidae | <i>Dolichopus</i> | spp. | 200 |
| Empididae | <i>Hydromia</i> | spp. | 82 |
| Ephydriidae | | spp. | 125 |
| Heleomyzidae | <i>Aecothea</i> | spp. | 2 |
| Heleomyzidae | <i>Heleomyza</i> | spp. | 380 |
| Heleomyzidae | <i>Nealeria</i> | spp. | 16 |
| Heleomyzidae | <i>Scoliocentra</i> | spp. | 36 |
| Heleomyzidae | | spp. | 86 |
| Milichiidae | | spp. | 2 |
| Mycetophilidae | <i>Mycetophila</i> | spp. | 250 |
| Mycetophilidae | | spp. | 185 |
| Piophilidae | | spp. | 11 |
| Psychodidae | | spp. | 1 |
| Scathophagidae | | spp. | 245 |
| Scatopsidae | | spp. | 3 |
| Sciaridae | | spp. | 145 |
| Sciomyzidae | | spp. | 1 |
| Simuliidae | | spp. | 640 |
| Sphaeroceridae | | spp. | 1 |
| Syrphidae | <i>Helophilus</i> | spp. | 14 |
| Tachinidae | <i>Exorista</i> | sp. 1 | 4 |
| Tachinidae | <i>Exorista</i> | sp. 2 | 3 |
| Tachinidae | <i>Exorista</i> | sp. 3 | 2 |
| Tachinidae | <i>Exorista</i> | sp. 4 | 7 |
| Tachinidae | <i>Exorista</i> | spp. | 22 |
| Tachinidae | <i>Nowickia</i> | spp. | 30 |
| Tachinidae | | spp. | 28 |
| Tipulidae | | spp. | 68 |
| Trichoceridae | | spp. | 14 |
| unsorted Greenland Material | | spp. | 1250 |